AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- 1 11. (CANCELED)
- (CURRENTLY AMENDED) A method for enhancing the sugar sucrose content
 and/or nutritional value ascorbic acid content of fruits of a plant of the genus
 Capsicum, the method comprising manipulating the CL and the Y loci which
 results in enhanced surose content and/or ascorbic acid content.
- (CURRENTLY AMENDED) The method according to claim 12, wherein the
 manipulation provides a plant of the genus Capsicum comprising two recessive y
 recessive y alleles and two recessive of alleles.
- 14. (CURRENTLY AMENDED) The method according to claim 13, wherein the y allele the y allele is derived obtained from a plant selected from the group consisting of Capsicum annuum, Capsicum baccatum, Capsicum frutescens, Capsicum chinense, and Capsicum chacoense.
- (CURRENTLY AMENDED) The method according to claim 13, wherein the y
 allele the y allele is derived obtained from Capsicum annuum.
- 16. (CURRENTLY AMENDED) The method according to claim 13, wherein the recessive el-allele c/ allele is derived obtained from a plant selected from the group consisting of apsicum annuum Capsicum annuum, Capsicum baccatum, Capsicum frutescens, Capsicum chinense, and Capsicum chacoense.
- (CURRENTLY AMENDED) The method according to claim 13, wherein the recessive el-allele cl allele is derived obtained from Capsicum annuum.
- 18. (CURRENTLY AMENDED) The method according to claim 12, wherein the manipulation results in a plant characterized by fruits having a <u>said</u> sucrose content which is at least <u>between</u> 1.5 times <u>and 3.4 times</u> higher than the sucrose content of <u>green immature</u> fruits of a plant of the genus <u>Capsicum of a</u>

- similar-type, said green immature fruits having at least one *CL* allele and at least one Y allele.
- (CURRENTLY AMENDED) The method according to claim 18, wherein the sucrose said sucrose content of the fruits is more than 5 between 5.4 grams and 6.2 grams per kilogram fresh weight.
- (CURRENTLY AMENDED) The method according to claim 18, wherein the sucrose said sucrose content of the fruits is 5-40 between 6.2 grams and 6.6 grams per kilogram fresh weight.
- (CURRENTLY AMENDED) The method according to claim 18, wherein the sucrose said sucrose content of the fruits is 5.4 to 16.8 between 6.6 grams and 7.1 grams per kilogram fresh weight.
- 22. (CANCELED)
- 23. (CURRENTLY AMENDED) The method according to elaim 22 claim 12, wherein the ascorbic acid content is at least between 1.3 times and 1.9 times higher than the ascorbic acid content in green immature fruits of a plant of the genus Capsicum of a similar type, said green immature fruits having at least one CL allele and at least one Y allele.
- (CURRENTLY AMENDED) The method according to claim 23, wherein <u>said</u>
 ascorbic acid content of the fruits is more than 2 <u>between 2.1</u> grams and 2.22
 grams per kilogram fresh weight.
- (CURRENTLY AMENDED) The method according to claim 23, wherein <u>said</u> ascorbic acid content of the fruits is 2-to-7 <u>between 2.22</u> grams <u>and 2.4 grams</u> per kilogram fresh weight.
- (CURRENTLY AMENDED) The method according to claim 23, wherein the wherein said ascorbic acid content of the fruits is 2.1 to 2.85 between 2.4 grams and 2.52 grams per kilogram fresh weight.
- 27.-28. (CANCELED)
- (CURRENTLY AMENDED) A method for increasing the sucrose content and the ascorbic acid content of fruits of a plant of the genus Capsicum, comprising

manipulating the CL and the Y loci to provide two recessive y-alleles y alleles and two recessive et alleles of alleles, wherein the sucrose content is increased to at least between 1.5 times and 3.4 times higher and wherein the ascorbic acid content is increased to between 1.3 times and 1.9 times higher than the sucrose content and the ascorbic acid content of green immature fruits of a plant of the genus Capsicum of a similar type, said green immature fruits having at least one CL allele and at least one Y allele.

- 30. (CANCELED)
- 31. (CURRENTLY AMENDED) The method of elaim 30 claim 29, wherein the sucrose content is increased to more than 5 between 5.4 grams and 7.1 grams per kilogram fresh weight, and the ascorbic acid content is increased to more than 2 between 2.1 grams and 2.52 grams per kilogram fresh weight.